Vol 17 No 3, March 2008
1674-1056/2008/17(03)/0825-04

Chinese Physics B

© 2008 Chin. Phys. Soc.
and IOP Publishing Ltd

Hawking radiation from the Schwarzschild black hole with a global monopole via gravitational anomaly*

Peng Jun-Jin(彭俊金) and Wu Shuang-Qing(吴双清)†

College of Physical Science and Technology, Central China Normal University, Wuhan 430079, *China*

(Received 11 July 2007; revised manuscript received 2 August 2007)

This paper derives the Hawking flux from the Schwarzschild black hole with a global monopole by using Robinson and Wilczek's method. Adopting a dimensional reduction technique, it can describe the effective quantum field in the $(3+1)$-dimensional global monopole background by an infinite collection of the $(1+1)$-dimensional massless fields if neglecting the ingoing modes near the horizon, where the gravitational anomaly can be cancelled by the $(1+1)$-dimensional black body radiation at the Hawking temperature.

Keywords: gravitational anomaly, Hawking radiation, black hole, global monopole
PACC: 0470, 1130J, 9760L

Recently, Robinson and Wilczek[1] (RW) proposed an intriguing approach to derive Hawking radiation from a Schwarzschild-type black hole through gravitational anomaly. Their basic idea goes as follows. Consider a massless scalar field in the higher dimensional space–time. Upon performing a dimensional reduction technique together with a partial wave decomposition, they found that the physics near the horizon in the original black hole background can be described by an infinite collection of the massless fields in a $(1+1)$-dimensional effective field theory. When omitting the classically irrelevant ingoing modes in the region near the horizon, the effective theory becomes chiral and exhibits gravitational anomaly in the near-horizon region, which just can be cancelled by the $(1+1)$-dimensional black body radiation at the Hawking temperature. As is shown later, the RW method is universal, and soon has been extended to other black hole cases[2−5] which contain gauge anomaly in addition to gravitational anomaly.

In this paper, we will use the RW method to investigate Hawking radiation of a static spherically symmetric Schwarzschild black hole with a global monopole from the viewpoint of cancelling the gravitational anomaly. During the process of the Grand Unified Theory (GUT) phase transition, we can imagine that a Schwarzschild black hole swallows a global monopole, forming a black-hole-global-monopole system. An unusual and stirring property of this composite system[6,7] is that it possesses a solid deficit angle, which makes it quite different topologically from that of a Schwarzschild black hole alone. Thermodynamical properties of such a static spherically symmetric system have been studied extensively in Ref.[7]. Because the background space–time considered here is not asymptotically flat rather it contains a topological defect due to the presence of a global monopole, the RW method, however, cannot be directly applied to derive the correct Hawking temperature of the black-hole-global-monopole system. In the following, we shall adopt a slightly different procedure and perform various coordinate transformations before we can use the RW method. Accordingly, we must generalize the RW method to the most general static case that the metric components satisfy $g_{tt}\cdot g_{rr}\neq -1$.

The metric of a Schwarzschild-type black hole with the global $O(3)$ monopole is described by[6,7]

$$\begin{aligned} \mathrm{d}s^2 &= -f(r)\mathrm{d}t^2 + f(r)^{-1}\mathrm{d}r^2 + r^2\mathrm{d}\Omega_2^2, \\ f(r) &= 1-\eta^2-2m/r, \end{aligned} \tag{1}$$

where m is the mass parameter of the black hole and η is related to the symmetry breaking scale when the global monopole is formed during the early universe. For a typical GUT symmetry breaking scale, $\eta^2\sim 10^{-6}$, so it is reasonable to assume $1-\eta^2\simeq 1$ throughout this paper.

Since the prime physical quantity obtained by means of the RW method is the Hawking temperature which enters into the first law of black hole thermodynamics, we begin with by reviewing the thermody-

*Project supported by the National Natural Science Foundation of China (Grant No 10675051).
†Corresponding author. E-mail: sqwu@phy.ccnu.edu.cn

namical properties of our black-hole-global-monopole system.

For the space–time metric (1), the Hawking temperature and the entropy are given by[7]

$$T = \frac{\kappa}{2\pi} = \frac{\partial_r f|_{r_{\rm H}}}{4\pi\sqrt{1-\eta^2}} = \frac{(1-\eta^2)^{3/2}}{8\pi m},$$
$$S = \frac{A}{4} = \pi r_{\rm H}^2 = \frac{4\pi m^2}{(1-\eta^2)^2}, \tag{2}$$

where κ and A are, respectively, the surface gravity and the area at the horizon $r_{\rm H} = 2m/(1-\eta^2)$.

The Arnowitt–Deser–Misner (ADM) mass M of the system can be calculated via the Komar integral

$$M = \frac{-1}{8\pi}\oint \xi^{\mu;\nu}_{(t)} {\rm d}^2\Sigma_{\mu\nu} = \frac{m}{\sqrt{1-\eta^2}}\,, \tag{3}$$

where $\xi^{\mu}_{(t)} = (1-\eta^2)^{-1/2}(\partial_t)^{\mu}$ is the normalized time-like Killing vector. Obviously, the ADM mass M is not equal to the mass parameter m because of the presence of a global monopole. One can easily show that the ADM mass M, the temperature T and the entropy S given above obey the differential and integral forms of the first law of black hole thermodynamics as follows

$${\rm d}M = T{\rm d}S, \quad M = 2TS. \tag{4}$$

Now introducing the following coordinate transformation

$$t \to (1-\eta^2)^{1/2}t\,, \quad r \to (1-\eta^2)^{-1/2}r\,, \tag{5}$$

and defining a new mass parameter

$$\widetilde{m} = (1-\eta^2)^{-3/2}m\,, \tag{6}$$

then we can rewrite the line element (1) as

$${\rm d}s^2 = -f(r){\rm d}t^2 + f(r)^{-1}{\rm d}r^2 + (1-\eta^2)r^2{\rm d}\Omega_2^2,$$
$$f(r) = 1 - 2\widetilde{m}/r\,. \tag{7}$$

This metric is, apart from the deficit solid angle $4\pi\eta^2$, very similar to the Schwarzschild solution. Remarkably, due to the presence of a global monopole, the original space–time Eq.(1) is not asymptotically flat but asymptotically bounded. After performing the above coordinate transformation, which is essentially a scale transformation, the metric is brought into an asymptotic one although it contains a topological defect. The advantage of this transformation is to make the calculation of the ADM mass and the Hawking temperature more feasible.

For the line element Eq.(7), the surface gravity at the horizon $r_{\rm H} = 2\widetilde{m}$ can be determined as $\kappa = \frac{1}{2}\partial_r f\big|_{r_{\rm H}}$. Analogously, one can compute the ADM mass $M = (1-\eta^2)\widetilde{m}$, the Hawking temperature $T = \kappa/(2\pi) = 1/(8\pi\widetilde{m})$, and the entropy $S = A/4 = 4\pi(1-\eta^2)\widetilde{m}^2$, and find that they are essentially identical to those given by Eqs.(2) and (3) by virtue of the relation Eq.(6), thus satisfying the same Bekenstein–Smarr relationship Eq.(4).

In the following, we will apply the RW method to show that the flux of Hawking radiation from a Schwarzschild-type black hole with the global monopole can be determined by anomaly cancellation conditions and regularity requirement at the horizon. The RW method, however, cannot be immediately applied to obtain the correct formula of Hawking temperature for the line element Eq.(1), rather it can be directly used to obtain that for the metric Eq.(7). Thus, we shall first base our analysis below upon the metric Eq.(7) but will soon turn to the space–time Eq.(1).

For simplicity, we consider the action for a massless scalar field in the background space–time Eq.(7). After performing the partial wave decomposition $\varphi = \sum_{lm}\varphi_{lm}(t,r)Y_{lm}(\theta,\phi)$, and only keeping the dominant terms, the action becomes

$$\begin{aligned} S[\varphi] &= -\frac{1}{2}\int {\rm d}^4x\sqrt{-g}g^{\mu\nu}\partial_\mu\varphi\partial_\nu\varphi \\ &= \frac{1}{2}\int {\rm d}t{\rm d}r{\rm d}\theta{\rm d}\phi(1-\eta^2)r^2\sin\theta\varphi\left[\frac{-1}{f}\partial_t^2\right. \\ &\quad \left. +\frac{1}{r^2}\partial_r\big(r^2 f\partial_r\big) + \frac{1}{(1-\eta^2)r^2}\Delta_\Omega\right]\varphi \\ &\approx \frac{1}{2}\sum_{lm}\int {\rm d}t{\rm d}r(1-\eta^2)r^2\ \varphi_{lm} \\ &\quad \times\left[\frac{-1}{f}\partial_t^2 + \partial_r\big(f\partial_r\big)\right]\varphi_{lm}\,, \end{aligned} \tag{8}$$

where Δ_Ω is the angular Laplace operator. Apparently, a free scalar field in the original $(3+1)$-dimensional background can be effectively described by an infinite collection of massless fields in the (1+1)-dimensional space–time with the effective metric

$${\rm d}s^2 = -f(r){\rm d}t^2 + f(r)^{-1}{\rm d}r^2, \tag{9}$$

together with the dilaton field $\Psi = (1-\eta^2)r^2$. On the other hand, if we start with the metric Eq.(1) and perform the dimension reduction, the same effective metric yields but with a different dilaton factor $\Psi = r^2$ whereas $f(r)$ is now given by Eq.(1).

Yet we can still go beyond further. For later usage, consider now the most general, static and spherically symmetric black hole solutions with the line element

$$\mathrm{d}s^2 = -f(r)\mathrm{d}t^2 + h(r)^{-1}\mathrm{d}r^2 + P(r)^2\mathrm{d}\Omega_2^2. \quad (10)$$

A similar dimension reduction technique leads to the $(1+1)$-dimensional effective metric

$$\mathrm{d}s^2 = -f(r)\mathrm{d}t^2 + h(r)^{-1}\mathrm{d}r^2 \quad (11)$$

with the dilaton field $\Psi = P(r)^2$, which makes no contribution to the anomaly. At the horizon $r = r_{\mathrm{H}}$, the surface gravity is $\kappa = \frac{1}{2}\sqrt{f'(r_{\mathrm{H}})h'(r_{\mathrm{H}})}$, where a prime denotes the derivative with respect to r.

Since we are considering a static background, the contribution from the dilaton field can be neglected. Thus we find that the physics near the horizon can be described by an infinite collection of massless fields in the $(1+1)$-dimensional effective theory on the background metric Eq.(11).

Next we turn to the gravitational anomaly. A gravitational anomaly is an anomaly in the general coordinate covariance, taking the form of nonconservation of the energy–momentum tensor. The consistent one arising in the $(1+1)$-dimensional chiral theory reads

$$\nabla_\mu T^\mu_{\ \nu} = \frac{1}{96\pi\sqrt{-g}}\varepsilon^{\beta\delta}\partial_\delta\partial_\alpha\Gamma^\alpha_{\nu\beta}, \quad (12)$$

on the other hand, the covariant anomaly for outgoing modes reads

$$\nabla_\mu \widetilde{T}^\mu_{\ \nu} = \frac{-1}{96\pi\sqrt{-g}}\varepsilon_{\mu\nu}\partial^\mu R\,, \quad (13)$$

where $\varepsilon^{\mu\nu}$ is an antisymmetric tensor with $\varepsilon^{tr} = 1$.

We will localize the physics outside the horizon since the effective theory is defined in the exterior region $[r_{\mathrm{H}}, +\infty]$. Now we divide the region outside the horizon into two parts: the near–horizon region $[r_{\mathrm{H}}, r_{\mathrm{H}} + \varepsilon]$, where the effective quantum field theory becomes chiral and exists a gravitational anomaly, and the other region $[r_{\mathrm{H}} + \varepsilon, +\infty]$, where the theory is not chiral and there is no anomaly. So let's focus on the anomaly in the region $[r_{\mathrm{H}}, r_{\mathrm{H}} + \varepsilon]$. Having omitted the classically irrelevant ingoing modes near the horizon, the energy–momentum tensor in this region exhibits an anomaly, which can be rewritten as

$$\nabla_\mu T^\mu_{(\mathrm{H})\nu} \equiv A_\nu \equiv \frac{1}{\sqrt{-g}}\partial_\mu N^\mu_{\ \nu}\,. \quad (14)$$

For a metric of the form as Eq.(11), $N^\mu_{\ \nu} = A_\nu = 0$ in the region $[r_{\mathrm{H}} + \varepsilon, +\infty]$. But in the near-horizon region $[r_{\mathrm{H}}, r_{\mathrm{H}} + \varepsilon]$, the components of $N^\mu_{\ \nu}$ are

$$\begin{aligned} N^t_t &= N^r_r = 0, \\ N^r_t &= \frac{1}{192\pi}\Big(f'h' + f''h\Big), \\ N^t_r &= \frac{-1}{192\pi h^2}\Big(h'^2 - h''h\Big). \end{aligned} \quad (15)$$

Taking into account the time independence of $T^\mu_{\ \nu}$, we can integrate Eq.(14), up to a trace $T^\alpha_\alpha(r)$, to get

$$\begin{aligned} T^t_t &= -\frac{K+Q}{f} - \frac{B(r)}{f} - \frac{I(r)}{2f} + T^\alpha_\alpha(r), \\ T^r_r &= \frac{K+Q}{f} + \frac{B(r)}{f} + \frac{I(r)}{2f}, \\ T^r_t &= -\sqrt{h/f}K + C(r) = -fhT^t_r\,, \end{aligned} \quad (16)$$

where $C(r) = \sqrt{h/f}\int_{r_{\mathrm{H}}}^r \sqrt{f(x)/h(x)}A_t(x)\mathrm{d}x$, $B(r) = \int_{r_{\mathrm{H}}}^r A_r(x)f(x)\mathrm{d}x$, $I(r) = \int_{r_{\mathrm{H}}}^r T^\alpha_\alpha(x)f'(x)\mathrm{d}x$, and K and Q are two integration constants. For the line element Eq.(11), $B(r)$ should be zero because $A_r = 0$ in the near-horizon region. In the limit $r \to r_{\mathrm{H}}$, we have $C(r) \to 0$, and $I(r)/f\big|_{r_{\mathrm{H}}} \to T^\alpha_\alpha(r_{\mathrm{H}})$.

Under the infinitesimal general coordinate transformations, the effective action varies as

$$\begin{aligned} & -\delta_\lambda W \\ &= \int \mathrm{d}^2x\sqrt{-g}\lambda^\nu\nabla_\mu T^\mu_{\ \nu} \\ &= \int \mathrm{d}t\mathrm{d}r\lambda^\nu\Big\{\partial_\mu\Big[N^\mu_\nu H(r)\Big] + \big[\sqrt{f/h}T^\mu_{(\mathrm{o})\nu} \\ &\quad -\sqrt{f/h}T^\mu_{(\mathrm{H})\nu} + N^\mu_\nu\big]\partial_\mu\Theta(r)\Big\} \\ &= \int \mathrm{d}t\mathrm{d}r\Big\{\lambda^t\Big(\partial_r\big[N^r_t H(r)\big] + \Big[N^r_t \\ &\quad +\sqrt{f/h}\big(T^r_{(\mathrm{o})t} - T^r_{(\mathrm{H})t}\big)\Big]\delta(r - r_{\mathrm{H}})\Big) \\ &\quad +\lambda^r\sqrt{f/h}\big(T^r_{(\mathrm{o})r} - T^r_{(\mathrm{H})r}\big)\delta(r - r_{\mathrm{H}})\Big\}\,, \end{aligned} \quad (17)$$

where $\Theta(r) = \Theta(r - r_{\mathrm{H}} - \varepsilon)$ is a scalar step function, $H(r) = 1 - \Theta(r)$ is a scalar top hat function, and we have written the total energy–momentum tensor T^μ_ν outside the horizon as

$$T^\mu_\nu = T^\mu_{(\mathrm{o})\nu}\Theta(r) + T^\mu_{(\mathrm{H})\nu}H(r)\,, \quad (18)$$

in which $T^\mu_{(\mathrm{o})\nu}$ is covariantly conserved and $T^\mu_{(\mathrm{H})\nu}$ obeys the anomalous Eq.(14). To derive the last expression of Eq.(17), we have taken the $\varepsilon \to 0$ limit.

In order to keep the diffeomorphism invariance, the variation of the effective action should vanish. The first term $\partial_r\big[N^r_t H(r)\big]$ in Eq.(17) can be cancelled by

the quantum effects of the ingoing modes. Setting $\delta_\lambda W = 0$, we obtain the following constraints

$$\left[\sqrt{f/h}\left(T^r_{(\mathrm{o})t} - T^r_{(\mathrm{H})t}\right) + N^r_t\right]\Big|_{r_\mathrm{H}} = 0,$$
$$\left(T^r_{(\mathrm{o})r} - T^r_{(\mathrm{H})r}\right)\Big|_{r_\mathrm{H}} = 0, \tag{19}$$

i.e.,

$$K_\mathrm{o} = K_\mathrm{H} + \Phi, \qquad Q_\mathrm{o} = Q_\mathrm{H} - \Phi, \tag{20}$$

where

$$\Phi = N^r_t(r_\mathrm{H}) = \frac{f'(r_\mathrm{H})h'(r_\mathrm{H})}{192\pi} = \frac{\kappa^2}{48\pi}\,. \tag{21}$$

In order to fix the four constants, we impose an additional regularity condition that requires the covariant energy–momentum tensor to vanish at the horizon. In the background of space–time Eq.(11), the covariant energy–momentum tensor is related to the consistent one by[8]

$$\sqrt{f/h}\widetilde{T}^r_t = \sqrt{f/h}T^r_t + \frac{h}{192\pi f}\left[ff'' - 2(f')^2\right]. \tag{22}$$

The condition $\widetilde{T}^r_t(r_\mathrm{H}) = 0$ yields $K_\mathrm{H} = -2\Phi$, which leads to $\sqrt{f/h}T^r_{(\mathrm{o})t}\big|_{r_\mathrm{H}} = -K_\mathrm{o} = \Phi$. So, Φ is the flux of Hawking radiation. A $(1+1)$-dimensional black body radiation has a flux of the form $\Phi = \dfrac{\pi}{12}T^2$, accurately giving the Hawking temperature $T = \kappa/(2\pi)$.

Applying the above analysis to the metric Eq.(7), we can obtain the correct Hawking temperature $T = 1/(8\pi\widetilde{m})$; whereas to the line element Eq.(1), we will get a different result $T = (1-\eta^2)^2/(8\pi m)$. So it is unadvisable to apply directly the RW method to the space–time Eq.(1). However, we can do the same analysis in another different way. By re-scaling $t \to \sqrt{1-\eta^2}\,t$, we rewrite the metric Eq.(1) as

$$\mathrm{d}s^2 = -f(r)\mathrm{d}t^2 + h(r)^{-1}\mathrm{d}r^2 + r^2\mathrm{d}\Omega_2^2,$$
$$f(r) = 1 - \frac{2m}{(1-\eta^2)r}, \quad h(r) = 1 - \eta^2 - \frac{2m}{r}, \tag{23}$$

and immediately derive the correct Hawking temperature $T = (1-\eta^2)^{3/2}/(8\pi m)$.

In summary, we have applied the RW method to derive the Hawking flux from a Schwarzschild-type black hole with the global monopole by requiring the cancellation of gravitational anomalies at the horizon. The flux has a form precisely equivalent to black body radiation with the Hawking temperature. To obtain the consistent expression of the Hawking temperature, it is not suitable to use the metric Eq.(1), otherwise one must divide the flux Φ by a factor $1-\eta^2$. Our analysis presented here can be directly applied to the case of a Schwarzschild–anti–de Sitter black hole with a global monopole[9] where $f(r) = 1-\eta^2-2m/r+r^2/l^2$.

References

[1] Robinson S P and Wilczek F 2005 *Phys. Rev. Lett.* **95** 011303

[2] Iso S, Umetsu H and Wilczek F 2006 *Phys. Rev. Lett.* **96** 151302
Iso S, Umetsu H and Wilczek F 2006 *Phys. Rev.* D **74** 044017
Murata K and Soda J 2006 *Phys. Rev.* D **74** 044018
Iso S, Morita H and Umetsu H 2007 *JHEP* **0704** 068

[3] Vagenas E C and Das S 2006 *JHEP* **0610** 025
Setare M R 2007 *Euro. Phys. J.* C **49** 865
Xu Z B and Chen B 2007 *Phys. Rev.* D **75** 024041
Xiao K, Liu W B and Zhang H B 2007 *Phys. Lett.* B **647** 482

[4] Jiang Q Q, Wu S Q and Cai X 2007 *Phys. Rev.* D **75** 064029
Jiang Q Q and Wu S Q 2007 *Phys. Lett.* B **647** 200
Jiang Q Q, Wu S Q and Cai X 2007 *Phys. Lett.* B **651** 48

[5] Shin H and Kim W 2007 *JHEP* **0706** 012

[6] Barriola M and Vilenkin A 1989 *Phys. Rev. Lett.* **63** 341

[7] Yu H W 1994 *Nucl. Phys.* B **430** 427

[8] Wu S Q and Peng J J 2007 *Class. Quantum Grav.* **24** 5123

[9] Chen J H, Jing J L and Wang Y J 2001 *Chin. Phys.* **10** 1071
Chen D Y, Jiang Q Q, Li H L and Yang S Z 2006 *Chin. Phys.* **15** 1425
Han Y W, Hong Y and Yang S Z 2007 *Acta Phys. Sin.* **56** 10 (in Chinese)